PHARMACY POLICY STATEMENT
North Carolina Marketplace

Crohn's Disease (CD)

For **initial** authorization:

1. Member is 18 years of age or older with moderately to severely active CD; AND
2. Medication must be prescribed by or in consultation with a gastroenterologist; AND
3. Must have had a negative TB test within the last 12 months; AND
4. Member has had a documented trial and inadequate response, or intolerance to at least **one** of the following conventional therapies: a 4-week trial of a corticosteroid OR a 12-week trial of 6-mercaptopurine, azathioprine, or methotrexate. Note: Trial is not required if member is switching from another biologic agent; OR
5. Member has severe disease that requires immediate use of a biologic agent, as indicated by one of the following:
   a) Extensive small bowel disease involving more than 100 cm;
   b) History of bowel or colon resection and is at high risk for CD recurrence (e.g., smoker, <30 years old, 2 or more resections, penetrating/fistulizing disease, etc.);
   c) Fistulizing disease.
6. **Dosage allowed/Quantity limit:**
   a) Induction (medical benefit): a one-time IV infusion based on weight. Up to 55 kg = 260 mg (2 vials); greater than 55 kg to 85 kg = 390 mg (3 vials); greater than 85 kg = 520 mg (4 vials);
   b) Maintenance (pharmacy or medical benefit): subcutaneous injection of 90 mg dose 8 weeks after induction and every 8 weeks thereafter.

Note to reviewer: A one-time induction dose is approved on the medical benefit. Maintenance therapy is

For **reauthorization**:
1. Chart notes have been provided showing improvement in signs and symptoms of CD (defined as mucosal healing, fewer flare-ups of symptoms, improved quality of life, etc.).

If all the above requirements are met, the medication will be approved for an additional 12 months.

Plaque Psoriasis (PsO)

For **initial** authorization:
1. Member must be 6 years of age or older; AND
2. Medication must be prescribed by or in consultation with a dermatologist; AND
3. Member has clinical documentation of moderate to severe plaque psoriasis characterized by 3% or more of body surface area (BSA) or disease affecting sensitive areas (e.g., hands, feet, face, genitals, etc.); AND
4. Must have had a negative TB test within the last 12 months; AND
5. Member has tried and failed to respond to treatment with at least **one** of the following:
   a) At least 12 weeks of photochemotherapy (i.e., psoralen plus ultraviolet A therapy);
   b) At least 12 weeks of phototherapy (i.e., UVB light therapy, Excimer laser treatments);
   c) At least a 4 week trial with topical antipsoriatic agents (i.e., anthralin, calcipotriene, coal tar, corticosteroids, tazarotene, tacrolimus, pimecrolimus); AND
6. Member has tried and failed, or unable to tolerate a systemic non-biologic DMARD (i.e., cyclosporine, methotrexate, acitretin) for at least 12 weeks.
7. **Dosage allowed/Quantity limit:**
   a) Adults: 100 kg or less: 45 mg subcutaneously at 0 and 4 weeks, and then every 12 weeks thereafter; more than 100 kg: 90 mg subcutaneously at 0 and 4 weeks, and then every 12 weeks thereafter;
   b) Pediatrics (6 to 17): subcutaneous dose by weight at week 0, week 4, and every 12 weeks thereafter. Less than 60 kg: 0.75 mg/kg; 60 kg to 100 kg: 45 mg; more than 100 kg: 90 mg.

If all the above requirements are met, the medication will be approved for 12 months.

For **reauthorization**:
1. Chart notes have been provided showing improvement of signs and symptoms of disease (e.g., documented member's BSA improvement, etc.).

If all the above requirements are met, the medication will be approved for an additional 12 months.

Psoriatic Arthritis (PsA)

For **initial** authorization:
1. Member must be 6 years of age or older; AND
2. Medication must be prescribed by or in consultation with a rheumatologist or a dermatologist; AND
3. Member has a documented diagnosis of active psoriatic arthritis (PsA); AND
4. Must have had a negative TB test within the last 12 months; AND
5. Member has met a 4-week trial of an NSAID taken at maximally tolerated doses AND a 3-month trial of a non-biologic DMARD agent (e.g., methotrexate, sulfasalazine, cyclosporine, etc.) unless **one** of the following situations is met:
   a) Non-biologic DMARD is not required for:

For **reauthorization**:

1. Chart notes have been provided showing improvement in signs and symptoms of UC (defined as clinical remission, decrease in rectal bleeding, decreased corticosteroid use, etc.).

If all the above requirements are met, the medication will be approved for an additional 12 months.

CareSource considers Stelara (ustekinumab) not medically necessary for the treatment of conditions that are not listed in this document. For any other indication, please refer to the Off-Label policy.

DATE	ACTION/DESCRIPTION
05/10/2017	Policy for Stelara created. Policies SRx-0042 and SRx-0043 archived. New diagnosis of Crohn's disease was added. For diagnosis of PsO: immunosuppressive criterion was separated from phototherapies and topical agents trials; TNF inhibitors Humira and Enbrel were listed as required trials; Psoriasis Area and Severity Index (PASI) score requirement was added. For diagnosis of PsA: TNF inhibitors Humira and Enbrel were listed as required trials. List of diagnoses considered not medically necessary was added.
11/13/2017	Age requirement for diagnosis of PsO updated.
02/26/2019	Humira was removed from criteria; Cimzia, Cosentyx, Otezla, Siliq and Xeljanz added to trial agents list. TB test allowed to be done within 12 months prior to initiation of therapy; chest x-ray option removed. Other drugs options allowed for PsA if there is an intolerance or contraindication to methotrexate.
10/31/2019	New indication of Ulcerative Colitis added.
11/23/2020	Removed repeat TB for reauth for all diagnoses. For CD: specified length of trials for conventional therapies, previously not specified. For severe disease, removed esophageal/gastroduodenal disease, specified that history of colonic resection must also be high risk for recurrence. Updated dosage [redacted]859 0.87

2. Sulz MC, Burri E, Michetti P, et al. Treatment Algorithms for Crohn's Disease. *Digestion*. 2020;101 Suppl 1:43-57.
3. Lichtenstein GR, Loftus EV, Isaacs KL, Regueiro MD, Gerson LB, Sands BE. ACG Clinical Guideline: Management of Crohn's Disease in Adults. *Am J Gastroenterol*. 2018;113(4):481-517.
4. Torres J, Bonovas S, Doherty G, et al. ECCO Guidelines on Therapeutics in Crohn's Disease: Medical Treatment. *J Crohns Colitis*. 2020;14(1):4-22.
5. Pimentel AM, Rocha R, Santana GO. Crohn's disease of esophagus, stomach and duodenum. *World J Gastrointest Pharmacol Ther*. 2019;10(2):35-49.
6. Terdiman JP, Gruss CB, Heidelbaugh JJ, Sultan S, Falck-Ytter YT; AGA Institute Clinical Practice and Quality Management Committee. American Gastroenterological Association Institute guideline on the use of thiopurines, methotrexate, and anti-TNF-α biologic drugs for the induction and maintenance of remission in inflammatory Crohn's disease. *Gastroenterology*. 2013;145(6):1459-1463.
7. Regueiro M, Velayos F, Greer JB, et al. American Gastroenterological Association Institute Technical Review on the Management of Crohn's Disease After Surgical Resection. *Gastroenterology*. 2017;152(1):277-295.e3.
8. Feagan BG, Sandborn WJ, Gasink C, et al. Ustekinumab as Induction and Maintenance Therapy for Crohn’s Disease. N Engl J Med 2016; 375:1946-1960.
9. Sandborn WJ, Feagan BG, Fedorak RN, et al. A randomized trial of Ustekinumab, a human interleukin-12/23 monoclonal antibody, in patients with moderate-to-severe Crohn's disease. *Gastroenterology*. 2008 Oct;135(4):1130-41. Epub 2008 Jul 17.
10.