
| MEDICAL POLICY STATEMENT Marketplace | | | | | |
|---------------------------------------|---------------------|--|--|--|--|
| Policy Name & Number | Date Effective | | | | |
| Facet Joint Interventions-MP-MM-1326 | -MM-1326 08/01/2024 | | | | |
| Policy Type | | | | | |

Facet Joint Interventions-MP-MM-1326 Effective Date: 08/01/2024



therapy, a physician supervised home exercise program (HEP), and/or chiropractic care.


HEP – A 6-week program requiring an exercise prescription and/or plan and a follow-up documented in the medical record after completion, or documentation of the inability to complete a HEP due to a stated physical reason (ie, increased pain, inability to physically perform exercises). Patient inconvenience or noncompliance without explanation does not constitute inability to complete.

 Inactive Conservative Therapies – Passive activities by the patient that aid in treating symptoms with pain including rest, ice, heat, medical devices, acupuncture, TENS use, and/or pharmacotherapy (prescription or over the counter [eq, NSAIDS, acetaminophen]).

Transcutaneous Electrical Nerve Stimulator (TENS) – A device that utilizes electrical current directed through electrodes placed on the surface of the skin to decrease the patient's perception of pain by inhibiting the transmission of afferent pain nerve impulses and/or stimulating the release of endorphins. Use, frequency, duration, and start dates must be documented in the medical record.

 Medial Branch Nerve Block Injection A diagnostic procedure in which a shortacting anesthetic (eg, lidocaine) is injected near small medial nerves connected to a specific facet joint. It may be performed to treat

Facet Joint Interventions-MP-MM-1326 Effective Date: 08/01/2024



- 4. Radiofrequency ablation is being considered as a therapeutic intervention.
- 5. Injections should be at least 2 weeks apart.
- 6. Imaging studies and physical exam ruled out other causes of spinal pain (eg, fracture, tumor, infection, herniated disk, spinal stenosis, significant deformity).
- 7. Patient history with at least 3 months of moderate to severe pain with functional impairment that has not adequately responded to active and inactive conservative therapy.
- 8. Failure of conservative therapy as evidenced by **ALL** of the following:
 - a. Documentation in the medical record of at least 6 weeks of active conservative therapy (as defined above) within the past 6 months OR inability to complete active conservative therapy due to contraindication, increased pain, or intolerance.
 - b. Documentation in the medical record of at least 6 weeks of inactive conservative therapy (as defined above) within the past six months.
- 9. No coagulopathy.
- 10. No current infection at the injection site.
- B. Diagnostic medial branch nerve blocks are **NOT** considered medically necessary when RFA is not being considered as a treatment option.
- II. Radiofrequency Ablation (RFA) for Facet Joint Pain
 - A. Initial RFA for facet joint pain is considered medically necessary when in 600.34 Tme

Facet Joint Interventions-MP-MM-1326 Effective Date: 08/01/2024