


MEDICAL POLICY STATEMENT

Marketplace

Policy Name & Number	Date Effective
Neonatal Discharge Criteria-MP-MM-1411	09/01/2024
Policy Type	
MEDIC(M)-3(TJ ET Q q 72.504 453.43 467.11 1!	

Indiana

Kentucky



A. Subject

Neonatal Discharge Criteria

B. Background

Infants who require neonatal admission remain at increased risk for morbidity and mortality following discharge. These infants require comprehensive discharge planning to ensure a smooth transition from the neonatal intensive care unit (NICU) and reduce morbidity and mortality after discharge.

Despite the inability to predict the exact timing of a NICU discharge, discharge planning should begin at NICU admission in an effort to avoid overwhelming parents and hospital staff. This planning will aid in minimizing discharge delays and will promote safe and healthy discharges to home.

Discharge may be appropriate when the establishment of physiologic competencies, including, but not limited to, thermoregulation, feeding, respiratory control, and stability regardless of weight or corrected gestational age, have been achieved.

C. Definitions

xminimizing discharge delays a6g06 465.75 Tm 0 g 0 G [<



10. Groothuis JR, Rosenberg AA. Home oxygen promotes weight gain in infants with bronchopulmonary dysplasia. *Am J Dis Child*. 1987;141(9):992-995. doi:10.1001/archpedi.1987.04460090069028
11. Halliday HL, Dumpit FM, Brady JP. Effects of inspired oxygen on echocardiographic assessment of pulmonary vascular resistance and myocardial contractility in bronchopulmonary dysplasia. *Pediatrics*. 1980;65(3):536-540. doi:10.1542/peds.65.3.536
12. Jefferies AL; Canadian Paediatric Society; Fetus and Newborn Committee. Going home: facilitating discharge of the preterm infant. *Paediatr Child Health*. 2014;19(1):31-42. doi:10.1093/pch/19.1.31
13. Kemper AR, Newman TB, Slaughter JL, et al. Clinical practice guideline revision: management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. *Pediatr*. 2022;150(3):e2022058859. doi:10.1542/peds.2022-058859
14. Moyer-Mileur LJ, Nielson DW, Pfeffer KD, et al. Eliminating sleep associated hypoxemia improves growth in infants with bronchopulmonary dysplasia. *Pediatr*. 1996;98(4 Pt 1):779-783. doi:10.1542/peds.98.4.779
15. Muchowski KE. Evaluation and treatment of neonatal hyperbilirubinemia. *Am Fam Physician*. 2014;89(11):873-878. Accessed February 26, 2024. www.aafp.org
16. Ortenstrand A, Waldenström U, Winbladh B. Early discharge of preterm infants needing limited special care, followed by domiciliary nursing care. *Acta Paediatr*. 1999;88(9):1024-1030. doi:10.1080/08035259950168568
17. Ortenstrand A, Winbladh B, Nordström G, et al. Early discharge of preterm infants followed by domiciliary nursing care: parents' anxiety, assessment of infant health and breastfeeding. *Acta Paediatr*. 2001;90(10):1190-1195. doi:10.1080/080352501317061639
18. Pinney MA, Cotton EK. Home management of bronchopulmonary dysplasia. *Pediatrics*. 1976;58(6):856-859. Accessed February 24, 2024. www.pubmed.ncbi.nlm.nih.gov
19. Schneiderman R, Kirkby S, Turenne W, et al. Incubator weaning in preterm infants and associated practice variation. *J Perinatol*. 2009;29(8):570-574. doi:10.1038/jp.2009.54
20. Sekar KC, Duke JC. Sleep apnea and hypoxemia in recently weaned premature infants with and without bronchopulmonary dysplasia. *Pediatr Pulmonol*. 1991;10(2):112-116. doi:10.1002/ppul.1950100213
21. Smith VC, Stewart J. Discharge planning for high-risk newborns. Accessed February 14, 2024. www.uptodate.com
22. Zecca E, Corsello M, Priolo F, et al. Early weaning from incubator and early discharge of preterm infants: randomized clinical trial. *Pediatr*. 2010;126(3):e651-e656. doi:10.1542/peds.2009-3005

Independent medical review #12/27/2021